161 Precalculus 1

**Review 2** 

In problems 1 – 4 consider the following quadratic function

$$y = -3x^2 + 7x + 1.$$

Problem 1 Write the function in the standard form

$$y = a(x - x_v)^2 + y_v$$

and find the coordinates of the vertex.

Solution. We first factor out the coefficient by  $x^2$ ,

$$y = -3\left(x^2 - \frac{7}{3}x - \frac{1}{3}\right).$$

Next we complete the square inside the parentheses using the formula

$$x^2 + bx = \left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2$$

In our case we have  $x^2 - \frac{1}{3}x = \left(x - \frac{7}{6}\right)^2 - \frac{49}{36}$ . After we plug in this

expression into the formula for y we get

$$y = -3\left(x - \frac{7}{6}\right)^2 + \frac{61}{12}$$

This is the standard form of the function  $y = -3x^2 + 7x + 1$ .

From the standard form we see that the coordinates of the vertex are

$$x_v = \frac{7}{6}, \quad y_v = \frac{61}{12}.$$

Problem 2 Find the x-intercepts of the graph of y (if any). Graph the function.

Solution. The *x*-intercepts can be found by quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{7 \pm \sqrt{7^2 - 4 \cdot (-3) \cdot 1}}{2 \cdot (-3)} = \frac{7 \pm \sqrt{61}}{-6}$$

The x-intercepts are located approximately at -0.13 and 2.47.

The graph of the function is shown below.



Problem 3 Consider function y over the restricted domain to the right from the vertex. Find the inverse function for this restriction of y. State the domain and the range of the inverse function.

Solution The restricted domain of y is the interval  $\left[\frac{7}{6},\infty\right]$ . To find an

expression for the inverse function we have to solve the equation  $y = -3x^2 + 7x + 1$  for x. Writing it as  $3x^2 - 7x + y - 1 = 0$  and applying the quadratic formula we get

$$x = \frac{7 \pm \sqrt{49 - 4 \cdot 3 \cdot (y - 1)}}{6} = \frac{7 \pm \sqrt{61 - 12y}}{6}$$

Because according to our choice of the restricted domain  $x \ge \frac{7}{6}$  the correct sign

in the above formula is +. Interchanging the input x and the output y we get the inverse function in the form

$$y = \frac{7 + \sqrt{61 - 12x}}{6}$$

Notice now that that the original function has the domain  $\left[\frac{7}{6},\infty\right)$  and the

range  $\left(-\infty, \frac{61}{12}\right]$ . Because for the inverse function we have to interchange the

domain and the range we see that the domain of the inverse function is

$$\left(-\infty,\frac{61}{12}\right]$$
 and its range is  $\left[\frac{7}{6},\infty\right)$ .

Problem 4. Graph the function y with the restricted domain and its inverse in the same coordinate system.

Solution Notice that the graph of y is the part of the graph from problem 2 to the right of the vertex. The graph of the inverse function can be obtained by interchanging the x and y coordinates, in other words by reflecting the graph of y about the line y = x. The graphs are shown below.



Problem 5 A penny is thrown up from the observation deck of the Empire State building (1050ft) with the velocity 50 ft/sec. Answer the following questions

- (a) After what time will the penny reach its maximum height?
- (b) What is the maximum height?
- (c) After what time will the penny hit the ground?

Solution. In a problem like this the height of the object after *t* seconds is given by the formula

$$h(t) = -16t^2 + v_0 t + h_0$$

where  $v_0$  is the initial velocity of the object in ft/sec and  $h_0$  is its initial height. In our case  $v_0 = 50$  and  $h_0 = 1050$  whence

$$h(t) = -16t^2 + 50t + 1050$$

Because the leading coefficient of the quadratic function h is negative the function takes it greatest value at the vertex.

(a) The *t* coordinate of the vertex is given by the formula

$$t_v = -\frac{b}{2a} = -\frac{50}{2 \cdot (-16)} = 1.5625 \, s$$

This answers question (a).

(b) The h coordinate of the vertex is given by

$$h_v = c - \frac{b^2}{4a} = 1050 - \frac{50^2}{4(-16)} = 1089.0625 \ ft$$

This answers question (b).

To answer question (c) we have to find the positive solution of the equation

$$h(t) = -16t^2 + 50t + 1050$$
$$-16t^2 + 50t + 1050 = 0$$

By the quadratic formula

$$t = \frac{-50 - \sqrt{50^2 - 4(-16)1050}}{2(-16)} \approx 9.812 \ s$$