


54. (a) A force diagram for one of the balls is shown on the right. The force of gravity  $m\vec{g}$  acts downward, the electrical force  $\vec{F}_e$  of the other ball acts to the left, and the tension in the thread acts along the thread, at the angle  $\theta$  to the vertical. The ball is in equilibrium, so its acceleration is zero. The  $y$  component of Newton's second law yields  $T \cos\theta - mg = 0$  and the  $x$  component yields  $T \sin\theta - F_e = 0$ . We solve the first equation for  $T$  and obtain  $T = mg/\cos\theta$ . We substitute the result into the second to obtain  $mg \tan\theta - F_e = 0$ .



Examination of the geometry of Figure 21-42 leads to

$$\tan\theta = \frac{x/2}{\sqrt{L^2 - (x/2)^2}}.$$

If  $L$  is much larger than  $x$  (which is the case if  $\theta$  is very small), we may neglect  $x/2$  in the denominator and write  $\tan\theta \approx x/2L$ . This is equivalent to approximating  $\tan\theta$  by  $\sin\theta$ . The magnitude of the electrical force of one ball on the other is

$$F_e = \frac{q^2}{4\pi\epsilon_0 x^2}$$

by Eq. 21-4. When these two expressions are used in the equation  $mg \tan\theta = F_e$ , we obtain

$$\frac{mgx}{2L} \approx \frac{1}{4\pi\epsilon_0} \frac{q^2}{x^2} \Rightarrow x \approx \left( \frac{q^2 L}{2\pi\epsilon_0 mg} \right)^{1/3}.$$

(b) We solve  $x^3 = 2kq^2L/mg$  for the charge (using Eq. 21-5):

$$q = \sqrt{\frac{mgx^3}{2kL}} = \sqrt{\frac{(0.010\text{kg})(9.8\text{m/s}^2)(0.050\text{m})^3}{2(8.99 \times 10^9 \text{N}\cdot\text{m}^2/\text{C}^2)(1.20\text{m})}} = \pm 2.4 \times 10^{-8} \text{C}.$$

Thus, the magnitude is  $|q| = 2.4 \times 10^{-8} \text{C}$ .