


1. On which intervals is the function $f(x) = -2x^3 + 3x^2 + 12x$ increasing? Where is it decreasing? Here is a picture from maple to help, but you must show your work.



Answer: Increasing over _____, decreasing over _____

2. Over what interval(s) is the above function concave up (leaning to the left) and concave down?

Answer: concave up over _____, concave down over _____.

3. Locate the maximum and minimum values of the function $f(x) = x^2 + \frac{2}{x}$ on the interval $[\frac{1}{2}, 2]$. Don't forget to check the endpoints as well as the critical points.

Answer: Minimum is _____, maximum is _____.

4. Approximate the cube root of 9 by using Newton's method to find the root of $y = x^3 - 9$. Start with the first guess of 2 and fill in the table.

x	y	y'
2	-1	?
?	?	

5. What does the mean value theorem say about the function $f(x) = x^3 - 9$ on the interval $[0, 3]$?

6. For question above, find the number in the interval $(0, 3)$ guaranteed by the mean value theorem to exist.

Use L'Hôpital's rule if applicable to find the following limits:

7. $\lim_{x \rightarrow 0} \frac{\cos(x) - 1}{x}$

8. $\lim_{x \rightarrow 0} \frac{x}{e^x}$

9. $\lim_{x \rightarrow 4} \frac{\sqrt{x} - 2}{x - 4}$

10. Definition: If f is continuous on $[a, b]$, the **definite integral of f from a to b** is

$$\int_a^b f(x) dx =$$

11. Define each symbol on the right hand side of the equal sign above.

12. Suppose $\int_a^b f(x) dx = -5$, $\int_a^c f(x) dx = 2$. What is $\int_c^b f(x) dx$?

13. For the function above, what is $\int_b^a f(x)dx$

14. Let $F(x) = \int_a^x f(t)dt$. F is a function of what variable? _____

15. For the function defined above, what is $F'(x)$?

16. Evaluate $\int_1^4 \frac{1}{\sqrt{t}} dt$

17. What are $\int_1^4 \frac{1}{\sqrt{x}} dx$ and $\int_1^4 \frac{1}{\sqrt{z}} dz$? Answer: _____ and _____

18. Let $F(x) = x \ln x - x$ show that $F'(x) = \ln x$ (Don't forget the product rule!)

19. Evaluate $\int_1^e \ln x dx$